Instruction for use Intensiv IPR Set

Selection of instruments for professional and calibrated application for the creation of interproximal space in IPR protocols.

Tray in stainless steel with silicone instrument holders. Inclusive content ready for sterilization.



Intensiv IPR Set. Ref. 6000/WG-69 LT Content of the stainless steel Tray: 1 Contra Angle, Ref. WG-69 LT, with light 1 Ejector, Ref. 053

1 Intensiv Ortho-Strips, Ref. OS08OP 1 Intensiv Ortho-Strips, Ref. OS80C 1 Intensiv Ortho-Strips, Ref. OS60 2 Intensiv Ortho-Strips, Ref. OS40

1 Intensiv Ortho-Strips, Ref. OS15 Intensiv ApproxOpener, Ref. AO2018DS
 Intensiv ApproxOpener, Ref. AO2018OS

1 Intensiv IPR-DistanceControl, Ref. IPR-DC010 1 Intensiv IPR-DistanceControl, Ref. IPR-DC015

 Intensiv IPR-DistanceControl, Ref. IPR-DC020
 Intensiv IPR-DistanceControl, Ref. IPR-DC040
 Intensiv IPR-DistanceControl, Ref. IPR-DC040 1 Intensiv IPR-DistanceControl, Ref. IPR-DC050

2 Intensiv ProxoContour Coarse, Ref. PXCC8040 2 Intensiv ProxoPolish, Ref. PXPO1508

Content of the stainless steel Tray:

1 Contra Angle, Ref. WG-69 A, without light 1 Intensiv Ortho-Strips, Ref. OS08OF

1 Intensiv Ortho-Strips Ref OS800 1 Intensiv Ortho-Strips, Ref. OS60 2 Intensiv Ortho-Strips, Ref. OS40

ntensiv IPR Set. Ref. 6000/WG-69 A

1 Intensiv Ortho-Strips, Ref. OS15 Intensiv ApproxOpener, Ref. AO2018DS 1 Intensiv ApproxOpener, Ref. AO2018OS 1 Intensiv IPR-DistanceControl, Ref. IPR-DC010

1 Intensiv IPR-DistanceControl, Ref. IPR-DC015

1 Intensiv IPR-DistanceControl, Ref. IPR-DC020 1 Intensiv IPR-DistanceControl, Ref. IPR-DC040 1 Intensiv IPR-DistanceControl, Ref. IPR-DC050

2 Intensiv ProxoContour Coarse, Ref. PXCC8040 2 Intensiv ProxoPolish, Ref. PXPO1508

WORKING SIDE

----->

Content of the stainless steel Trave 1 Intensiv Ortho-Strips Ref OS08OF 1 Intensiv Ortho-Strips, Ref. OS80C 1 Intensiv ApproxOpener, Ref. AO2018DS



Mechanical instruments for the creation of interproximal space

Intensiv Ortho-Strips

ersity of Zurich

Oscillating instruments for the interdental reduction of

To be used in combination with the contra-angles:

- Intensiv Swingle, WG-69 LT Ortho, with light
 Intensiv Swingle, WG-69 A Ortho, without light
- KaVo head 61 LG
- KaVo contra-angle E61C

Product description

- Double sided diamond-coated oscillating metal strip Strip height: 3.7 mm
 Length of the diamond-coated part: 13 mm
- Sterilizable and reusable
- Opening of the interdental space

Contouring and polishing of the enamel

Recommendations for use

- Apply Intensiv Ortho-Strips in interproximal areas according to the IPR enamel reduction protocol for or thodontic treatments.
- A pre-opening of tight contact points using Intensive Ortho-Strips Opener is suggested, Ref. OS08OP/6, Ref. OS08OP-R/6, right or Ref. OS08OP-L/6, left.
- · In case of very tight contact points separate the teet by placing a wooden wedge cervically. This makes stripping more comfortable both for operator and pa-
- Set the micro motor at 40'000 RPM/min max. With Intensiv Swingle this corresponds to 20'000 oscillations per minute approx. (2:1).
- Activate water spray during the entire treatment nimum 50 ml /min) Regulate water flow in case of water excess or short-
- Insert the strip in a cervical direction while the contra-
- Contact pressure: bend the strip 1-1.5 mm to achieve
- the ideal 1-2.5N operating force.

 Do not stop the strip in the interdental space
- Remove the strip from the interdental space whilst the contra-angle is moving

Intensiv Ortho-Strips Coarse

Oscillating instruments for a fast and efficient reduction of interproximal enamel in orthodontic treatments

To be used in combination with the contra-angles

- KaVo head 61 LG
- KaVo contra-angle E61C

Product description

- Oscillating diamond-coated and perforated metal strip
 Diamond grit-size: 80 µm
- side of the strip, thickness 0.2±0.05 mm

 One-Sided Right (R): diamond coating on the right
- side of the strip, thickness 0.2±0.05 mm
- Double-Sided: diamond coating on both sides of the strip, thickness0.3±0.05 mm
- Strip height: 3.7 mm
- Length of the working part: 13 mm
 Sterilizable and reusable

during orthodontic IPR protocol.

Recommendations for use

- Apply Intensiv Ortho-Strips Coarse in interproximal areas according to the IPR enamel reduction protocol for orthodontic treatments.
- In case of very tight contact points separate the teeth by placing a wooden wedge cervically. This makes tripping less uncomfortable both for operator and
- Activate water spray during the entire treatment (minimum 50 ml/min)

- the ideal 1-2.5N operating force.
 Do not stop the strip in the interdental space.
- Remove the strip from the interdental space whilst the contra-angle is moving

tight interdental contact points

• KaVo contra-angle E61C

Product description

- Serrated oscillating diamond-coated metal strip
- Available in 3 versions
- of the strip
 One-Sided Right (R): diamond coating on the right
- side of the strip
- Strip height: 3.7 mm

- Sterilizable and reusable

stage of the orthodontic IPR procedure.

- areas at the beginning of the IPR orthodontics proto-col for the opening of tight contact points. Subsequently proceed with the enamel reduction pro-
- tocol using Intensiv Ortho-Strips System.
- tensiv Swingle this corresponds to 20'000 oscillations
- (minimum 50 mL/min).
- Regulate water flow in case of water excess or shortage

Unilateral or bilateral interproximal enamel reduction

- A pre-opening of tight contact points using Intensi
- Ortho-Strips Opener is suggested, Ref. OS08OP/6, Ref. OS08OP-R/6, right or Ref. OS08OP-L/6, left.
- Set the micro motor at 40'000 RPM/min max. With In-
- tensiv Swingle this corresponds to 20'000 oscillations per minute approx. (2:1).
- Regulate water flow in case of water excess or short-
- Insert the strip in a cervical direction while the contraangle is moving. This prevents the instruments from
- Contact pressure: bend the strip 1-1.5 mm to achieve

Intensiv Ortho-Strips Opener

Oscillating instruments for easy and rapid opening of

To be used in combination with the contra-angles Intensiv Swingle, WG-69 LT Ortho, with light
 Intensiv Swingle, WG-69 A Ortho, without light

KaVo head 61 LG

- Diamond grit-size: 8 µm
- One-Sided Left (L): diamond coating on the left side
- Double-Sided: diamond coating on both sides of the
- Length of the working part: 13 mm Strip thickness: 0.1±0.03 mm

Opening of tight interdental contact areas at the initial

- Recommendations for use
 Apply Intensiv Ortho-Strips Opener in interproximal
- In case of very tight contact points separate the teeth by placing a wooden wedge cervically. Set the micro motor at 40'000 RPM/min max. With In-
- per minute approx. (2:1).

 Activate water spray during the entire treatment

achieve the ideal 1-2.5N operating force. Do not stop the strip in the interdental space Remove the strip from the interdental space whilst the

Risk Warning for all mechanical instruments

- cause breakage of the instrument.

 Avoid temperatures above 180°C which may affect the
- cause damage to the instrument.

 To ensure traceability of the instruments during their entire application, it is recommended to keep the

Maintenance and Sterilization of all mechanical

- instruments
 Instruments are delivered non sterile. They must be sterilized prior to first use and disinfected, cleaned, and sterilized after each use
- Clean the instruments and remove debris after each use, so as to maintain their abrasive properties. Use a single-pulsed or fractionated vacuum autoclave
- and subvacuum drying. Chemiclave sterilizers may also be used. Hot air sterilizers are not suitable for diamond instruments. Sterilization must be carried out according to valid
- dated procedures.
 Sterilization process is referred to ISO standard 17664.
- It is recommended to follow the indications below Cycles at 134°C: Tmin = 134°C – Tmax = 138°C Pressure = 3.15 bar abs
- Time = 4 min (raisable) Tmin = 121°C - Tmax = 125°C Pressure = 2.10 bar abs

Packaging of 3, 6, 12 Intensiv Ortho-Strips, Ref. OS90 Intensiv Ortho-Strips, Ref. OS60 Intensiv Ortho-Strips Ref OS40

Intensiv Ortho-Strips, Ref. OS15 Intensiv Ortho-Strips Set. Ref. 060A (OS90, OS40, OS25,

Intensiv Ortho-Strips Set, Ref. 060B (OS60, OS40, OS25,

Intensiv Ortho-Strips Coarse, Double-Sided, Ref. OS80C Intensiv Ortho-Strips Coarse, One-Sided, Ref. OS80C-R.

Intensiv Ortho-Strips Coarse, One-Sided, Ref. OS80C-L.

Intensiv Ortho-Strips Opener, Double-Sided, Ref OS08OP siv Ortho-Strips Opener, One-Sided,

Intensiv Ortho-Strips Opener, One-Sided,

• Insert the strip in a cervical direction while the contra-Intensiv ApproxOpener angle is moving. This prevents the instrument from Product description • Manual metal strip

Operative pressure: bend the strip 1-1.5 mm to

contra-angle is moving

- Avoid bending and levering: use of excessive force (strong levering effect with the contra-angle) could
- lead to the breakage of the instrument.

 Avoid jamming or levering actions when oscillating, as this increases the risk of instrument breakage.
- Oscillating over 20'000 oscillations per minute could
- durability of the instrument. Avoid applying forces greater than 3N as this could

- Intensiv ApproxOpener is positioned incisally/coro nally between the teeth in accordance with the inter-dental space and moved in gingival direction.

 Depending on the initial positioning of Intensive Ap
 - proxOpener in the interdental space, the forward or return movement will be active or passive

- Serrated and diamond coated (8µm) on one side - Serrated and diamond coated (8µm) on both sides

Overall length: 20mm
 Thickness of the serrated and one-side diamond

coated strip: approx. 0.09mm

Thickness of the serrated and two-side diamond

Opening of tight interproximal areas in orthodontic treatments (one-side and two-side diamond coated

coated strip: approx. 0.10mm

Total height of the instrument: 18mm

Total length of the instrument: 27 5mm

Removal of black triangles

Versions

Height: 4mm

- Risk Warnings It is recommended not to bend the working part or use it as a lever, applying a force superior than 2N, in order to prevent a break of the instrument.
- Please pay attention to the force applied to the instru-ment when working in the vicinity of soft tissue; apply a force similar to that used with a manual finishing strip to prevent mucosal lesions. Intensiv ApproxOpener is held firmly between fingers and secured with dental floss.

Metal strip serrated and diamond coated on one side Ref. AO2018OS/3 Package of 3 Strips

Metal strip serrated and diamond coated on both sides Ref. AO2018DS/3 Package of 3 Strips

Intensiv ProxoContour Coarse

Manual instruments for the creation of interproximal space

Product description

- Metal strip, diamond-coated on one side only with two working parts of different grit sizes: (80µm, blue, 40um red)
- One sided, 3 rows of perforation, diamond-coated
- metal strip Perforated anatomical ends for a better gripping
- Height: 2.5mm
 Total length: 80mm

Thickness: 0.05mm Sterilizable and reusable

· Unilateral interproximal enamel reduction and finishing during orthodontic IPR protocol (Stripping).

- The total length of the strip (80mm) allows an optimal finger support on the adjacent teeth during work
- To get past the contact area, use the uncoated part of the strip and exert a slight horizontal movement. Do not twist, bend or pull strip too tightly during use.

 • After introduction, work with the less abrasive side
- more abrasive side (80µm, blue).
 Then apply the 80µm diamond-coated side, with cautious horizontal movements, to model the proximal contours. Next, switch again to the less abrasive side (40µm, red) to complete the procedure with the fin
- After finishing to perform the polishing of the surfaces, it is recommended to use Intensiv ProxoPolish with a grain grit of 15µm and 8µm for the polishing. Risk Warnings Intensiv Intensiv ProxoContour Coarse
- Bending or twisting the metal strip during use can lead to fractures and/or fissures. Pulling too forcefully in tight interdental spaces or beginning with the more abrasive (80µm) grit can cause the fissures.
- Bending or twisting the metal strip can lead to detachment of the diamond coating.

 • Leaving proximal surfaces roughened with 40µm grain size can lead to undesirable mechanical retention and

Intensiv ProxoPolish Pat. 699 819

formation of plaque

- Product description
 Metal strip, with only one diamond-coated side and two working parts in different grain sizes, 15 µm (yellow) and 8 µm (orange)

 Perforated anatomical ends for a better gripping
- Height: 2.5mm
- Total length: 80mm
 Thickness: 0.05mm

· Sterilizable and reusable Unilateral interproximal enamel polishing during orthodontic IPR protocol (Stripping)


1 Intensiv IPR-DistanceControl Ref IPR-DC010

Intensiv IPR-DistanceControl, Ref. IPR-DC010
Intensiv IPR-DistanceControl, Ref. IPR-DC020
Intensiv ProxoContour Coarse, Ref. PXCC8040
Intensiv ProxoPolish, Ref. PXPO1508

The total length of the strip (80mm) allows an ideal support of fingers on adjacent teeth during treat-

After introducing the strip, apply a back and forth

- movement, first with the slightly more abrasive side (15µm, yellow).
 To complete the treatment of surfaces with polishing,
- finish with less abrasive side (8µm, orange).



ment of the diamond coating

The strip is supplied non-sterile. Prior to first use on

the natient and immediately after each use, the instru


- ents must be disinfected, cleaned, and sterilized. Keep the strip separate from non-stainless steel instru ments, such as rubber polishers and abrasive stones. If disinfection by immersion is being used, it is impor-
- tant to use only cleaning/disinfection solutions that provide corrosion protection and strictly observe the concentrations recommended by the manufacturer. (40um, red), in order to facilitate the access of the Sterilization must be performed with validated proce dures. If possible, use a single-pulsed or fractionated vacuum autoclave and subvacuum drying. Chemiclave
 - sterilizers may also be used. Sterilization process is referred to ISO standard 17664. nded to follow the indications bel
 - Cycles at 134°C: Tmin = 134°C Tmax = 138°C Pressure = 3.15 bar abs Time = 4 min (raisable) Cycles at 121°C:
 - Pressure = 2.10 bar abs Time = 16 min (raisable) (grit size 80um and 40um)

(grit size 15um and 8um)

Tmin = 121°C - Tmax = 125°C

Ref. PXCC8040/6 Package of 6 Strips

Ref. PXPO1508/6 Package of 6 Strips



TWIST

- interdental space with the thickness gauge immediately below or through a combination of 2 or more in-If the designated instrument has too much play in the

 Intensiv IPR-DistanceControl is supplied non-sterile.
 Sterilize the instruments before first use. After each use on patients, the instruments must be disinfected cleaned and sterilized as well. Use a single-pulsed or fractionated vacuum autoclave

Risk Warnings During insertion pay attention to the soft tissue to avoid injuries.

Packaging Kit containing 8 elements, Ref. IPR-DC Set Pack containing 3 instruments of the same thickness

DON'T TWIST

tions for use contained in Intensiv IPR Set packaging

www.wh.com

Ignaz-Glaser-Strasse 53 5111 Bürmoos Austria office@wh.com






Tel. +41 91 986 50 50 Fax +41 91 986 50 59 info@intensiv.ch www.intensiv.ch

Measurement of the created proximal space

and subvacuum drying. Chemiclave sterilizers may also


Ref IPR-DC010/3- IPR-DC015/3- IPR-DC020/3 IPR-DC025/3; IPR-DC030/3; IPR-DC040/3; IPR-DC050/3;

Reciprocating Contra Angle () 0297 **Intensiv Swingle**





Intensiv Swingle, WG-69 LT Ortho, with light Intensiv Swingle, WG-69 A Ortho, without light



IPR-DistanceControl

- is a manual metal instrument
- 0,20 mm
- 0.40 mm
- 5 metallic elements of different thicknesses with er gonomic grip: - 0.10 mm

Intensiv IPR (InterProximalReduction)-DistanceControl

- - If the designated instrument does not enter into the interdental space created, proceed to measure the

· Height of measuring part: 4mm

- Measurement and control of the teeth distance in the various stages of the IPR technique in orthodontics.
- Length of measuring part: 40mm
 Total length of instrument: 50mm Sterilizable and reusable

• IPR-DC is inserted in the vestibular/buccal direction.

interdental space created, proceed to measure the in-terdental space with the thickness gauge immediately above or through a combination of 2 or more instru-

Sterilization must be carried out according to vali-